

AN EXTENSION OF BIRAN'S LAGRANGIAN BARRIER THEOREM

GUANG-CUN LU

(Communicated by Jon Wolfson)

ABSTRACT. We use the Gromov-Witten invariants and a nonsqueezing theorem by the author to affirm a conjecture by P.Biran on the Lagrangian barriers.

1. MAIN RESULTS

A Kähler manifold is a triple consisting of a symplectic manifold (M, ω) and an integrable complex structure J compatible with ω on M . If $[\omega] \in H^2(M^{2n}, \mathbb{Z})$ it follows from Kodaira's embedding theorem that there exists a smooth and reduced complex hypersurface $\Sigma \subset M$ such that its homology class $[\Sigma] \in H_{2n-2}(M)$ represents the Poincaré dual $k[\omega] \in H^2(M)$ for some $k \in \mathbb{N}$. Following [1] $\mathcal{P} = (M, \omega, J; \Sigma)$ is called a *smoothly polarized Kähler manifold*. Under the conditions that either $\dim_{\mathbb{R}} M \leq 6$ or $\omega|_{\pi_2(M)} = 0$ the following two theorems were proved in Theorem 1.D and Theorem 4.A of [1] respectively.

Theorem 1.1. *If (M, ω) is a Kähler manifold with $[\omega] \in H^2(M, \mathbb{Q})$, then for every $\epsilon > 0$ there exists a Lagrangian CW-complex $\Delta_{\epsilon} \subset (M, \omega)$ such that every symplectic embedding $\varphi : B(\epsilon) \rightarrow (M, \omega)$ must satisfy $\varphi(B(\epsilon)) \cap \Delta_{\epsilon} \neq \emptyset$.*

Theorem 1.2. *If $\mathcal{P} = (M, \omega, J; \Sigma)$ is a n -dimensional polarized Kähler manifold of degree k then every symplectic embedding $\varphi : B^{2n}(\lambda) := \{x \in \mathbb{R}^{2n} \mid |x| \leq \lambda^2\} \rightarrow (M, \omega)$ with $\lambda^2 \geq \frac{1}{k\pi}$ must intersect the skeleton $\Delta_{\mathcal{P}}$ associated to the polarization $\mathcal{P}$.*

For the definitions of Δ_{ϵ} and $\Delta_{\mathcal{P}}$ the reader may refer to [1]. Actually, such generalizations were conjectured in Remark 4.B of [1]. As in [1] Theorem 1.1 may be derived from Theorem 1.2.

Following [2] a Stein manifold is said to be *subcritical* if it admits a plurisubharmonic function which has only critical points of index less than half the real dimension. A polarization $\mathcal{P} = (M, \omega, J; \Sigma)$ is called *subcritical* if the complement $(M \setminus \Sigma, \omega)$ is a subcritical Stein manifold. From the proof of Theorem 1.2 we easily get the following generalizations of Theorem F and Theorem G in [2].

Date: July 1, 2003 and, in revised form, December 26, 2003.

2000 Mathematics Subject Classification. Primary 57R17, 53D35, 53D40; Secondary 32Q15, 32Q28.

Key words and phrases. Polarized Kähler manifolds, Lagrangian barrier, Gromov-Witten invariants, nonsqueezing theorem, Gromov width.

The author was supported in part by NNSF 19971045 and 10371007 of China.

Theorem 1.3. *If a closed Kähler manifold (M, ω, J) admits a subcritical polarization $\mathcal{P} = (M, \omega, J; \Sigma)$ of degree k , then $\mathcal{W}_G(M, \omega) \leq \frac{1}{k}$ and $k \leq \dim_{\mathbb{C}} M$. Moreover, if the linear system of holomorphic sections of the normal bundle to Σ , $N_{\Sigma/M} \rightarrow \Sigma$, is base point free then $\mathcal{W}_G(M, \omega) = \frac{1}{k} \geq 1/\dim_{\mathbb{C}} M$. Here $\mathcal{W}_G(M, \omega)$ stands for the Gromov width of (M, ω) .*

2. PROOF OF THEOREMS

Our purpose is to prove Theorem 1.2. The ideas are similar to those of Biran. However, we use the theory of virtual cycles and some techniques in [4] to compute the desired the Gromov-Witten invariant without additional assumptions as in [1]. Then we directly use the author's previous work on pseudo symplectic capacities and avoid Gromov's arguments as used in [1]. For convenience of the reader we need to recall some related notions in [1]. A subset Δ of a symplectic manifold (M, ω) is called an *embedded CW-complex* if there exists an abstract finite and path-connected CW-complex K and a homeomorphism $i : K \rightarrow \Delta \subset M$ such that for every cell $C \subset K$ the restriction $i|_{\text{Int}(C)} : \text{Int}(C) \rightarrow M$ is a smooth embedding. When the image $i(\text{Int}(C))$ by i of each cell C in K is an isotropic submanifold of (M, ω) the above embedded CW-complex Δ is called *isotropic*. The dimension $\dim \Delta$ of Δ is defined as the maximum of those of cells in K . An embedded isotropic CW-complex of dimension $\frac{1}{2} \dim_{\mathbb{R}} M$ in (M, ω) is called a *Lagrangian CW-complex*.

Another key role in [1] is the *standard symplectic disc bundle*. For a closed symplectic manifold (S, σ) with $[\sigma] \in H^2(S; \mathbb{Z})$ there exists a Hermitian line bundle $p : L \rightarrow S$ with $c_1(L) = [\sigma]$ and a compatible connection ∇ on L with curvature $R^\nabla = 2\pi i \sigma$ (cf. [5, Prop. 8.3.1]). Denote by $\|\cdot\|$ the Hermitian metric and by $E_L = \{v \in L \mid \|v\| < 1\}$ the open unit disc bundle of L . Let α^∇ the associated *transgression 1-form* on $L \setminus 0$ with $d\alpha^\nabla = -p^*\sigma$, and r the radial coordinate along the fibres induced by $\|\cdot\|$. It was shown that

$$\omega_{\text{can}} := p^*\sigma + d(r^2\alpha^\nabla)$$

is a symplectic form on E_L and that the symplectic type of $(E_L, \omega_{\text{can}})$ depends only on the symplectic type of (S, σ) and the topological type of the complex line bundle $p : L \rightarrow S$. Moreover, $(E_L, \omega_{\text{can}})$ is uniquely characterized (up to symplectomorphism) by the following three properties:

- All fibres of $p : E_L \rightarrow S$ are symplectic with respect to ω_{can} and have area 1.
- The restriction of ω_{can} to the zero section $S \subset E_L$ equals σ .
- ω_{can} is S^1 -invariant with respect to the obvious circle action on E_L .

Following [1] we call

$$p : (E_L, \omega_{\text{can}}) \rightarrow (S, \sigma)$$

a *standard symplectic disc bundle over (S, σ) modelled on L* and $p : (E_L, c\omega_{\text{can}}) \rightarrow (S, \sigma)$ a *standard symplectic disc bundle with fibres of area c* for each $c > 0$.

As done in [1] this standard symplectic disc bundle can be compactified into a $\mathbb{C}P^1$ -bundle over (S, σ) . Indeed, let $\mathbb{C}$ stand for the trivial complex line bundle over S . Then the direct sum $L \oplus \mathbb{C}$ is a complex vector bundle of rank 2 over S . Denote by $p_X : X_L = P(L \oplus \mathbb{C}) \rightarrow S$ its projective bundle, which is a $\mathbb{C}P^1$ -bundle over S . It has two distinguished sections: the zero section $Z_0 = P(0 \oplus \mathbb{C})$ and the section at infinity $Z_\infty = P(L \oplus 0)$. Clearly, the open manifold $X_L \setminus Z_\infty$ is diffeomorphic to the disc bundle E_L . Let F_s stand for the fibre of X_L at $s \in S$. It was shown

in [1] that a given $J_S \in \mathcal{J}(S, \sigma)$ and the connection ∇ determine a unique almost complex structure J_L on the total space of L . This J_L induces the almost complex structures J_E on E and J_X on X_L again. In particular, $J_E \in \mathcal{J}(E_L, \omega_{\text{can}})$ and Z_0, Z_∞ and all fibres F_s of $p_X : X_L \rightarrow S$ are holomorphic with respect to J_X . These show that $p_X : (X_L, J_X) \rightarrow (S, J_S)$ is an almost complex fibration with fibre $(\mathbb{CP}^1, i)$ in the following sense.

Definition 2.1 ([1, Def.6.3.A]). Let (F, J_F) , (B, J_B) and (X, J) be three almost complex manifolds. A holomorphic map $p : (X, J) \rightarrow (B, J_B)$ is called an almost complex fibration with fibre (F, J_F) if every $b \in B$ has a neighborhood U and a trivialization $\varphi : X|_U \rightarrow U \times F$ such that for every $a \in U$ the map $\varphi|_{F_a} : (F_a, J|_{F_a}) \rightarrow (a \times F, J_F)$ is holomorphic.

For every $0 < \rho < 1$ both $E_L(\rho) := \{v \in E_L \mid \|v\| < \rho\}$ and $\overline{E_L(\rho)} := \{v \in E_L \mid \|v\| \leq \rho\}$ are subbundles of E_L .

Lemma 2.2 ([1, Lem. 5.2.A]). *Let $p : (E_L, \omega_{\text{can}}) \rightarrow (S, \sigma)$ be a standard symplectic disc bundle modeled on a Hermitian line bundle $p : L \rightarrow S$ as above. Then there exists a diffeomorphism $f : E_L \rightarrow X_L \setminus Z_\infty$ and a family of symplectic forms $\{\eta_\rho\}_{0 < \rho < 1}$ on X_L such that*

- (1) $f^*\eta_\rho = \omega_{\text{can}}$ on $\overline{E_L(\rho)}$ for every $0 < \rho < 1$;
- (2) f sends the fibres of $E_L \rightarrow S$ to the fibres of $X_L \setminus Z_\infty \rightarrow S$;
- (3) If S is identified with the zero-section of E_L then $f(S) = Z_0$ and $p_X \circ f|_S : S \rightarrow S$ is the identity map;
- (4) Z_0, Z_∞ and all fibres $F_s = p_X^{-1}(s)$ ($s \in S$) of $p_X : X_L \rightarrow S$ are not only symplectic with respect to η_ρ for every $0 < \rho < 1$, but also holomorphic with respect to J_X . Moreover, for every $0 < \rho < 1$ it holds that $\eta_\rho|_{TZ_0} = (p_X^*\sigma)|_{TZ_0}$ and $\eta_\rho|_{TZ_\infty} = c_\rho(p_X^*\sigma)|_{TZ_\infty}$ for some $0 < c_\rho < 1 - \rho^2$;
- (5) The area of the fibres F_s satisfies $\rho^2 < \int_{F_s} \eta_\rho < 1$ for every $0 < \rho < 1$.

Moreover, if $J_S \in \mathcal{J}(S, \sigma)$ then $J_X \in \mathcal{J}(X_L, \eta_\rho)$ for every $0 < \rho < 1$. In particular, if (S, σ, J_S) is Kähler and $L \rightarrow S$ is a holomorphic line bundle then (X_L, η_ρ, J_X) is Kähler for every $0 < \rho < 1$.

It was proved in Lemma 6.A of [1] that the almost complex structure J_X is regular for the class of the fibre $F \in H_2(X_L, \mathbb{Z})$ and that the space of J_X -holomorphic spheres in the class F is made up of exactly the fibres $F_s = p_X^{-1}(s)$, $s \in S$. So the space $\overline{M}_{0,3}(X_L, J_X, F)$ of all J_X -holomorphic stable maps in class F of genus 0 and with 3 marked points is nonempty. As expected we have:

Lemma 2.3. *Let $F \in H_2(X_L; \mathbb{Z})$ denote the homology class of a fibre of $X_L \rightarrow S$. Then the Gromov-Witten invariant of (X_L, η_ρ) ,*

$$\Psi_{F,0,3}^{(X_L, \eta_\rho)}(pt; [Z_0], [Z_\infty], pt) = 1.$$

That is, (X_L, η_ρ) is a strong 0-symplectic uniruled manifold in the sense of Definition 1.16 in [3].

Proof. Let $A = [\mathbb{CP}^1] \in H_2(\mathbb{CP}^1; \mathbb{Z})$ and $i : \mathbb{CP}^1 \rightarrow X_L$ be the inclusion. Then $F = i_*(A)$. Since the Gromov-Witten invariants are symplectic deformation invariants, we may fix a $\eta \equiv \eta_\rho$. By Lemma 2.2 all fibres $F_s = \pi_X^{-1}(s)$ ($s \in S$) are not only

symplectic with respect to η but also holomorphic with respect to J_X . Thus for each $s \in S$ and $x \in F_s$ the tangent space $T_x F_s$ is a symplectic and $J_X(x)$ -invariant subspace of $(T_x X_L, \eta_x)$. Let H_x denote the η_x -orthogonal complement of $T_x F_s$. Then $T_x X_L = H_x \oplus T_x F_s$ and $Dp_X(x)|_{H_x} : H_x \rightarrow T_s S$ is an isomorphism. Note that J_X is also compatible with η . We have that $\eta(J_X \xi, J_X \eta) = \eta(\xi, \eta)$ for all $\xi, \eta \in TX_L$. This implies that H_x is also a $J_X(x)$ -invariant subspace of $T_x X_L$. Consequently, $J_X(x)$ preserve the splitting $T_x X_L = H_x \oplus T_x F_s$. Moreover, the projection $p_X : (X_L, J_X) \rightarrow (S, J_S)$ is holomorphic (cf. [1, §6.3]). Hence the almost complex structure J_X on X_L is *fibered* in the sense of Definition 2.2 in [4].

Since any stable map has connected image set it is easily checked that each stable map in $\overline{M}_{0,3}(X_L, J_X, F)$ must also entirely lie in a fibre of X_L . It follows from this and Lemma 4.3 in [4] that for any representative $\tilde{\tau} = (\Sigma, \tilde{h})$ of $\tau \in \overline{M}_{0,3}(X_L, J_X, F)$ the cokernel of $D\tilde{\partial}_{J_X}(\tilde{h})$ can be spanned by elements of the space

$$\mathcal{L}_{\tilde{h}}^V = L^{1,p} \left(\wedge^{0,1} (\tilde{h}^*(T_{vert} X_L)) \right),$$

where $V = T_{vert} X_L$ is the vertical tangent bundle of TX_L . By Proposition 4.3 in [4] one can choose R and the embeddings e in the construction of the virtual moduli cycle $\overline{M}_{0,3}^\nu(X_L, J_X, F)$ of $\overline{M}_{0,3}(X_L, J_X, F)$ such that $e_{\tilde{\tau}}(\nu) \in \mathcal{L}_{\tilde{h}}^V$ for all $\nu \in R$ and all $\tilde{\tau}$. That is, we can choose a fibered pair (J_X, ν) on X_L in the sense of Definition 4.4 in [4]. As proved in Proposition 4.4 of [4] such a virtual moduli cycle $\overline{M}_{0,3}^\nu(X_L, J_X, F)$ has the following property. For each element of it, a parameterized stable map $(\Sigma, \tilde{h})$, each component $\tilde{h}_k$ of $\tilde{h}$ satisfies an equation of the form $\tilde{\partial}_{J_X} \tilde{h}_k = \nu_k$ for some section ν_k of $\wedge^{0,1}(\tilde{h}_k^*(T_{vert} X_L))$. This implies that each $p_X \circ \tilde{h}_k : S^2 \rightarrow S$ is (i, J_S) -holomorphic and therefore that each element in $\overline{M}_{0,3}^\nu(X_L, J_X, F)$ has the image contained in a single fibre of X_L . For any $s \in S$ we identify $(F_s, J_X|_{F_s})$ with $(\mathbb{C}P^1, i)$. Then the subset $\overline{M}_{0,3}^\nu(X_L, J_X, F)_s$ of $\overline{M}_{0,3}^\nu(X_L, J_X, F)$ consisting of stable maps with image in F_s regularizes the space $\overline{M}_{0,3}(\mathbb{C}P^1, i, A)$ in the sense of Definition 4.3 in [4]. Hence

$$\Psi_{F,0,3}^{(X_L, \eta)}(pt; [Z_0], [Z_\infty], pt) = \Psi_{A,0,3}^{(\mathbb{C}P^1, \omega_{FS})}(pt; pt, pt, pt) = 1.$$

Here we have used pt to stand for the point classes in the different spaces. Lemma 2.3 is proved. $\square$

Recall that for a closed symplectic manifold (M, ω) and homology classes $\alpha_0, \alpha_\infty \in H_*(M, \mathbb{Q})$ we in [3, Def.1.8] defined a number

$$\text{GW}_g(M, \omega; \alpha_0, \alpha_\infty) \in (0, +\infty]$$

by the infimum of the ω -areas $\omega(A)$ of the homology classes $A \in H_2(M; \mathbb{Z})$ for which the Gromov-Witten invariant $\Psi_{A,g,m+2}(C; \alpha_0, \alpha_\infty, \beta_1, \dots, \beta_m) \neq 0$ for some homology classes $\beta_1, \dots, \beta_m \in H_*(M; \mathbb{Q})$ and $C \in H_*(\overline{\mathcal{M}}_{g,m+2}; \mathbb{Q})$ and integer $m > 0$. Moreover, in Definition 1.25 of [3] we also defined another number $\text{GW}(M, \omega) \in (0, +\infty]$ by

$$\text{GW}(M, \omega) = \inf \text{GW}_g(M, \omega; pt, \alpha),$$

where the infimum is taken over all nonnegative integers g and all homology classes $\alpha \in H_*(M; \mathbb{Q}) \setminus \{0\}$ of degree $\deg \alpha \leq \dim M - 1$. Theorem 1.26 in [3] claimed that

$$(1) \quad \mathcal{W}_G(M, \omega) \leq \text{GW}(M, \omega)$$

for any symplectic uniruled manifold (M, ω) .

Proof of Theorem 1.2. By the decomposition Theorem 2.6.A in [1], $(M \setminus \Delta_{\mathcal{P}}, \omega)$ is symplectomorphic to the standard symplectic disc bundle $(E_{N_{\Sigma}}, \frac{1}{k}\omega_{\text{can}}) \rightarrow (\Sigma, \sigma = k\omega|_{\Sigma})$ over Σ which is modeled on the normal bundle N_{Σ} and with fibres of area $1/k$. Assume that $\varphi(B(\lambda)) \cap \Delta_{\mathcal{P}} = \emptyset$ for some symplectic embedding $\varphi : B^{2n}(\lambda) \rightarrow (M, \omega)$ with $\lambda^2 \geq \frac{1}{k\pi}$. Then there exists a symplectic embedding $\psi : B(\lambda) \rightarrow (E_{N_{\Sigma}}, \frac{1}{k}\omega_{\text{can}})$. Since $\psi(B(\lambda))$ is compact there exists a positive number $\rho \in (0, 1)$ such that $\psi(B(\lambda)) \subset E_{N_{\Sigma}}(\rho)$. Lemma 2.2 gives a symplectic embedding from $\psi(B(\lambda))$ into $(X_{N_{\Sigma}}, \frac{1}{k}\eta_{\rho})$. Hence

$$(2) \quad \mathcal{W}_G(X_{N_{\Sigma}}, \frac{1}{k}\eta_{\rho}) \geq \pi\lambda^2 \geq 1/k.$$

On the another hand, by the definition of $\text{GW}(M, \omega)$ and Lemmas 2.2, 2.3 we have

$$\text{GW}(X_{N_{\Sigma}}, \eta_{\rho}) \leq \int_{F_s} \eta_{\rho} < 1$$

for this ρ . This and (1) together give

$$\mathcal{W}_G(X_{N_{\Sigma}}, \frac{1}{k}\eta_{\rho}) \leq \text{GW}(X_{N_{\Sigma}}, \frac{1}{k}\eta_{\rho}) \leq \frac{1}{k} \int_{F_s} \eta_{\rho} < \frac{1}{k},$$

which contradicts to (2). Theorem 1.2 is proved. $\square$

Acknowledgements. I am grateful to Kai Cieliebak for sending me some preprints. I would like to thank the referees for several advice for improving the presentation, pointing out a careless mistake in the statement of Theorem 1.3 in the previous version of the paper.

REFERENCES

- [1] P. Biran, *Lagrangian barriers and symplectic embeddings*, Geom. funct. anal. **11** (2001), 4070-464.
- [2] P. Biran and K. Cieliebak, *Symplectic topology on subcritical manifolds*, Comment. Math. Helvetici **76** (2001), 712-753.
- [3] G. C. Lu, *Gromov-Witten invariants and pseudo symplectic capacities*, math.SG/0103195, v7, May 21, 2004.
- [4] D. McDuff, *Quantum homology of fibrations over S^2* , International Journal of Mathematics, **11** (2000), 665-721.
- [5] N. M. J. Woodhouse, *Geometric Quantization*, Second edition, Oxford University Press, New York, 1991.

DEPARTMENT OF MATHEMATICS, BEIJING NORMAL UNIVERSITY, BEIJING 100875, P.R.CHINA
E-mail address: gclu@bnu.edu.cn